

*PHNIP Country Health Statistical Report*

# Zambia

---

**September 2002**



---

Population, Health and Nutrition Information Project  
600 13th Street, NW, Suite 710, Washington, DC 20005  
E-mail: [dolphnhelp@phnip.com](mailto:dolphnhelp@phnip.com)

# Table of Contents

---

## 1. About the Report

## 2. Country Profile

### I. Statistical Overview

- *Demographic Indicators*
- *Socioeconomic Indicators*
- *Family Planning Indicators*
- *Maternal Health Indicators*
- *Child Survival Indicators*
- *HIV/AIDS Prevention Indicators*
- *Infectious Diseases Control Indicators*

### II. Trends in Health Indicators

- *Current and Projected Population by Age and Sex*  
(Figure 1 and Table 1.1)
- *Population Estimates and Projections*  
(Table 1.2)
- *Estimated Total Fertility Rate per Woman*  
(Figure 2 and Table 2)
- *Infant Mortality Rates / Under-5 Mortality Rates*  
(Figure 3 and Table 3)
- *Vaccination Coverage (DPT3) Trends*  
(Figure 4 and Table 4)
- *Contraceptive Prevalence Rates - Married Women*  
(Figure 5 and Table 5)

## 3. Data Notes

## 4. Sources

# About the Report

Released September 2002

---

This report is one of a series of Country Health Statistical Reports produced by the Population, Health and Nutrition Information (PHNI) Project. Each profile contains statistical data on current health conditions, population dynamics, health and family planning behavior, and health and population trends in a given developing country. Information is compiled from PHNI's health statistics database, which draws data from a diverse range of sources listed at the end of this profile.

Hard copy editions of PHNI's profiles are available from our publications department. Reports will soon be available on the USAID Global Health Web site in Portable Document Format (.pdf). Any feedback you have on the content or presentation of this report would be greatly appreciated. We would also appreciate receiving any more recent, more accurate, or more representative information.

Contact us at the following:



Population, Health and Nutrition Information Project  
600 13th Street, NW, Suite 710  
Washington, DC 20005

Phone: 202-393-9001  
Fax: 202-393-9019  
E-mail: [dolphnhelp@phnip.com](mailto:dolphnhelp@phnip.com)

The PHNI Project, a USAID project managed by Jorge Scientific Corporation, with The Futures Group International and John Snow, Inc., prepared this document under the Data for Decision Making Project (936-3098.01), contract number HRN-C-00-00-00004-00, with the Bureau for Global Health, U.S. Agency for International Development (USAID).

# I. Statistical Overview

\*see data notes

Demographic Indicators				
Indicator	Value	Data Unit	Year	Source
Total Population	9,770,199		2001	BUCEN - IDB-2000
Population Growth Rate	1.9	%	2001	BUCEN - IDB-2000
Percent Urban	39.5	%	1999	World Bank/WDI-2001
Women, 15-19	624,935		2002	BUCEN - IDB-2000
Women, 15-49	2,251,616		2002	BUCEN - IDB-2000
Life Expectancy at Birth	37.3		2001	BUCEN - IDB-2000
Crude Birth Rate	41.5	per 1,000	2001	BUCEN - IDB-2000
Crude Death Rate	22	per 1,000	2001	BUCEN - IDB-2000
Number of Live Births	405	000s	2001	BUCEN - IDB-2000

Socioeconomic Indicators				
Indicator	Value	Data Unit	Year	Source
GNI per Capita (PPP)	720	\$	1999	World Bank/WDI-2001
Health Expenditure as Percentage of GDP	6.9	%	1990-1998	World Bank/WDI-2001
Physicians per 1,000 People	0.09		1990-1998	World Bank/WDI-2001
Adult Literacy Rate	77.2	%	1999	World Bank/WDI-2001
Adult Literacy Rate, Female	70.2	%	1999	World Bank/WDI-2001
Adult Literacy Rate, Male	84.6	%	1999	World Bank/WDI-2001
Gross Enrollment Rate - Primary School	101	%	1998	UNESCO-2000
Gender Parity Index - Net Enrollment Rate	1		1998	UNESCO-2000
Access to an Improved Water Source (Rural)	48	%	2000	World Bank/WDI-2001
Access to an Improved Water Source (Urban)	88	%	2000	World Bank/WDI-2001
Access to Improved Sanitation Facilities (Rural)	64	%	2000	World Bank/WDI-2001
Access to Improved Sanitation Facilities (Urban)	99	%	2000	World Bank/WDI-2001

Family Planning Indicators				
Indicator	Value	Data Unit	Year	Source
Total Fertility Rate (BUCEN)	5.5		2001	BUCEN - IDB-2000
Total Fertility Rate (DHS)	6.1		1996	Zambia DHS -1996
Contraceptive Prevalence Rate, Modern Methods, All Women	11.2	%	1996	Zambia DHS -1996
Contraceptive Prevalence Rate, Modern Methods, Married Women	14.4	%	1996	Zambia DHS -1996
Median Age of Sexual Debut Among Women, Ages 25-49	16.3		1996	Zambia DHS -1996
Mean Ideal Family Size	5.3		1996	Zambia DHS -1996
Women 20-24 Who Gave Birth Before Age 20	63.4	%	1996	Zambia DHS -1996

Maternal Health Indicators				
Indicator	Value	Data Unit	Year	Source
Maternal Mortality Ratio (DHS)	649	Per 100,000 live births	1990-1996	Zambia DHS -1996
Maternal Mortality Ratio (WHO/Hill)	867	Per 100,000 live births	1995	WHO/Hill-2001
Antenatal Care (at least 1 visit)	94	%	1996	Zambia DHS -1996
Antenatal Care (2+ visits)	92.1	%	1996	Zambia DHS -1996
Antenatal Care (4+ visits)	69.9	%	1996	Zambia DHS -1996
Assisted Delivery by a Health Professional	46.5	%	1996	Zambia DHS -1996
Assisted Delivery by Doctor	3.6	%	1996	Zambia DHS -1996
Assisted Delivery by Other Health Professional	42.9	%	1996	Zambia DHS -1996

Child Survival Indicators				
Indicator	Value	Data Unit	Year	Source
<b>Acute Respiratory Infection (ARI) and Oral Rehydration Therapy (ORT)</b>				
ARI Treatment - Children Under 5	70.7	%	1996	Zambia DHS -1996
ORT Use Rate (ORS, RHS, or Increased Fluids)	75.7	%	1996	Zambia DHS -1996
<b>Mortality Indicators</b>				
Infant Mortality Rate (BUCEN)	89	per 1,000 live births	2002	BUCEN - IDB-2000
Infant Mortality Rate (DHS)	108.9	per 1,000 live births	1996	Zambia DHS -1996
Infant Mortality Rate, Females (BUCEN)	83	per 1,000 live births	2002	BUCEN - IDB-2000
Infant Mortality Rate, Males (BUCEN)	96	per 1,000 live births	2002	BUCEN - IDB-2000
Under-5 Mortality Rate (BUCEN)	166.4	per 1,000 live births	2001	BUCEN - IDB-2000
Under-5 Mortality Rate (DHS)	196.6	per 1,000 live births	1996	Zambia DHS -1996
<b>Nutrition Indicators</b>				
Exclusive Breastfeeding (under 4 mos.)	26.3	%	1996	Zambia DHS -1996
Exclusive Breastfeeding (under 6 mos.)	19.3	%	1996	Zambia DHS -1996
Stunted (height-for-age)	42.4	%	1996	Zambia DHS -1996
Underweight (weight-for-age)	23.5	%	1996	Zambia DHS -1996
Wasted (weight-for-height)	4.2	%	1996	Zambia DHS -1996
<b>Vaccination Coverage</b>				
DPT3 Vaccination Rate (DHS)	85.7	%	1996	Zambia DHS -1996
Measles Vaccination Rate (DHS)	86.5	%	1996	Zambia DHS -1996
Polio Vaccination Rate (DHS)	84.4	%	1996	Zambia DHS -1996
Tetanus Toxoid Vaccination	35.4	%	1996	Zambia DHS -1996

HIV/AIDS Prevention Indicators				
Indicator	Value	Data Unit	Year	Source
HIV Prevalence, Adults	19.9	%	1999	UNAIDS-2000
HIV Prevalence, Females 15-24, High Estimates	18.7	%	1999	UNAIDS-2000
HIV Prevalence, Females 15-24, Low Estimates	16.9	%	1999	UNAIDS-2000
HIV Prevalence, Males 15-24, High Estimates	9.3	%	1999	UNAIDS-2000
HIV Prevalence, Males 15-24, Low Estimates	7.1	%	1999	UNAIDS-2000
Males Reporting Condom Use With Last Non-Regular Partner	3.5 <sup>1</sup>	%	1996	Zambia DHS -1996
Females Reporting Condom Use With Last Non-Regular Partner	2.1 <sup>2</sup>	%	1996	Zambia DHS -1996


Infectious Diseases Control Indicators				
Indicator	Value	Data Unit	Year	Source
TB Incidence Rate	Data Not Available			
TB Case Detection Rate	Data Not Available			
TB (DOTS) Treatment Success Rate	Data Not Available			
Malaria	3,215,866	Reported # of cases	1996	WHO/RBM-2002

<sup>1</sup> The value for this country represents the percentage of males reporting use of condom at last sexual encounter with any person (cohabitating or not) other than the man's spouse.

<sup>2</sup> The value for this country represents the percentage of females reporting use of condom at last sexual encounter with any person (cohabitating or not) other than the woman's spouse.

## II. Trends in Health Indicators

**Figure 1: Current and Projected Population by Age and Sex (in thousands)**



**Table 1.1: Current and Projected Population by Age and Sex**

Source: BUCEN - IDB

Age Group	Male - 2000	Male - 2020	Female - 2000	Female - 2020
0-4	863,172	994,082	851,202	975,752
5-9	747,396	933,143	742,259	920,584
10-14	679,991	893,377	677,484	884,347
15-19	598,464	839,920	592,163	829,900
20-24	497,534	758,312	477,887	740,013
25-29	398,863	638,483	367,260	607,585
30-34	284,248	509,175	265,720	466,583
35-39	192,017	372,857	190,379	327,946
40-44	120,149	255,280	138,881	219,222
45-49	88,288	176,073	115,250	157,316
50-54	71,542	115,691	100,726	121,566
55-59	62,427	77,171	89,077	99,673
60-64	55,785	49,504	75,727	81,990
65-69	42,461	37,125	57,732	70,951
70-74	29,212	29,159	36,993	58,533
75-79	19,568	21,829	22,543	43,789
80+	14,202	19,838	15,816	37,855


**TABLE 1.2: Population Estimates and Projections**

Source: BUCEN - IDB

1980	1990	2000	2010	2020
5,699,777	7,850,881	9,582,418	11,482,431	13,364,624



**Figure 2: Estimated Total Fertility Rate per Woman**




**Table 2: Estimated Total Fertility Rate per Woman**

Source: BUCEN - IDB

1980	1985	1990	1995	2000	2005	2010	2015	2020	2025
7.2	6.8	6.5	6.1	5.6	5.2	4.7	4.2	3.8	3.5

**Figure 3: Infant Mortality Rates / Under-5 Mortality Rates**




**Table 3: Infant Mortality Rates / Under-5 Mortality Rates**

Source: BUCEN - IDB 2000

Indicator	1980	1985	1990	1995	2000	2005	2010	2015	2020
Infant Mortality Rate (BUCEN)	89.4	87.6	106.0	99.1	92.4	84.9	77.9	70.6	62.8
Under-5 Mortality Rate (BUCEN)	165.5	162.1	190.6	178.8	168.8	156.7	145.7	133.5	119.3

**Figure 4: Vaccination Coverage (DPT3) Trends**




**Table 4: Vaccination Coverage (DPT3) Trends**

Source: WHO/Global Summary

1990	1995	2000
71.0	82.0	75.5

**Figure 5: Contraceptive Prevalence Rates, Modern Methods, Married Women**



**Table 5: Contraceptive Prevalence Rates, Modern Methods, Married Women**

Source: Zambia DHS

1992	1996
8.9	14.4

# Data Notes

---

\*Indicator definitions that differ from those in the data notes section are identified by footnotes accompanying the indicator tables.

## **Demographic Indicators**

<b>Total Population</b>	The number of people in a given area (i.e., country) in a particular time period (usually a midyear estimate).
<b>Population Growth Rate</b>	The average annual growth rate is the rate of natural increase in a population plus the net migration rate. The rate of natural increase is the difference between the birth rate and the death rate, but it is conventionally measured in percentage terms (per hundred rather than per thousand).
<b>Percent Urban</b>	The percentage of the midyear population living in areas defined as urban in each country and reported to the United Nations.
<b>Women, 15-19</b>	The number of women between the ages of 15-19 in the midyear population.
<b>Women, 15-49</b>	The number of women between the ages of 15-49 in the midyear population.
<b>Life Expectancy at Birth</b>	The average number of years that a person at age 0 will live if age-specific death rates remain constant. Life expectancy at birth is highly affected by rates of infant and child death.
<b>Crude Birth Rate</b>	The number of births per thousand of the population. The product of the number of live births, divided by the midpoint population, multiplied by 1,000.
<b>Crude Death Rate</b>	The number of deaths per thousand of the population. The product of the number of deaths divided by the midpoint population, multiplied by 1,000.
<b>Number of Live Births</b>	The number of live births, annually, within a country.

## **Socioeconomic Indicators**

<b>GNI per Capita (PPP)</b>	The gross national income (GNI) converted to international dollars using purchasing power parity (PPP) rates. An international dollar has the same purchasing power over GNI as a U.S. dollar has in the United States.
<b>Health Expenditure as Percentage of GDP</b>	Total health expenditures is the sum of public and private health expenditures figured as a percentage of a country's gross domestic product (GDP). It covers the provision of health services (preventive and curative), family planning activities, nutrition activities, and emergency aid designated for health but does not include provision of water and sanitation.
<b>Physicians per 1,000 People</b>	Physicians are defined as graduates of any faculty or school of medicine who are working in the country in any medical field (practice, teaching, research).
<b>Adult Literacy Rate</b>	The percentage of people ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.
<b>Adult Literacy Rate, Female</b>	The percentage of women ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.
<b>Adult Literacy Rate, Male</b>	The percentage of men ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.
<b>Gross Enrollment Rate - Primary School</b>	Total enrolment in primary level of education, regardless of age, expressed as a percentage of the official school-age population corresponding to primary level of education in give school-year. GER can be over 100% due to the inclusion of over-aged and under-aged pupils/students because of early or late entrants, and grade repetition.

<b>Gender Parity Index - Net Enrollment Rate</b>	The ratio of the female-to-male net primary school enrollment rates measures progress towards gender equity in school attendance and the level of learning opportunities available for women in relation to those available to men. Calculation Method: Divide the female net primary school enrollment rate by the male net primary school enrollment rate.
<b>Access to an Improved Water Source (Rural)</b>	Access to an improved water source in rural areas. Refers to the percentage of the rural population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs. Reasonable access is defined as the availability of at least 20 liters a person a day from a source within one kilometer of the dwelling.
<b>Access to an Improved Water Source (Urban)</b>	Access to an improved water source in urban areas. Refers to the percentage of the urban population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs. Reasonable access is defined as the availability of at least 20 liters a person a day from a source within one kilometer of the dwelling.
<b>Access to Improved Sanitation Facilities (Rural)</b>	Access to improved sanitation facilities in rural areas. Refers to the percentage of the rural population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.
<b>Access to Improved Sanitation Facilities (Urban)</b>	Access to improved sanitation facilities in urban areas. Refers to the percentage of the urban population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.
<b>Family Planning Indicators</b>	
<b>Total Fertility Rate (BUCEN)</b>	The number of children a woman between ages 15-44 would have during her reproductive life, if, for all of her childbearing years she were to experience the age-specific birth rates for that given year.
<b>Total Fertility Rate (DHS)</b>	The number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.
<b>Contraceptive Prevalence Rate, Modern Methods, All Women</b>	Percentage of all women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).
<b>Contraceptive Prevalence Rate, Modern Methods, Married Women</b>	Percent of currently married ages women 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).
<b>Median Age of Sexual Debut Among Women, Ages 25-49</b>	Median age of first sexual intercourse for women ages 25-49.
<b>Mean Ideal Family Size</b>	Mean ideal number of children for all women, according to number of living children.
<b>Women 20-24 Who Gave Birth Before Age 20</b>	Percentage of women ages 20-24 who have already given birth or were pregnant with their first child before the age of 20.
<b>Maternal Health Indicators</b>	
<b>Maternal Mortality Ratio (DHS)</b>	The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births, arrived mostly through 'direct sisterhood method'. The use of information reported by a sibling is to expand the sample size and to compensate for the lack of vital registration system.
<b>Maternal Mortality Ratio (WHO/Hill)</b>	The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births arrived through Hill et al., adjustment procedure depending on the nature of data used.
<b>Antenatal Care (at least 1 visit)</b>	Percentage of women of reproductive age (15-49) who receive at least one antenatal care visit during pregnancy (in the three-year period preceding the survey).
<b>Antenatal Care (2+ visits)</b>	Percentage of women of reproductive age (15-49) who receive at least two antenatal care visits during pregnancy (in the three-year period preceding the survey).

<b>Antenatal Care (4+ visits)</b>	Percentage of women of reproductive age (15-49) who receive four or more antenatal care visits during pregnancy (in the three-year period preceding the survey).
<b>Assisted Delivery by a Health Professional</b>	The percentage of births/deliveries that occur with the assistance of any trained health professional during the five-year period preceding the survey. May include doctors, nurses, midwives, village health workers, or any other trained health professional.
<b>Assisted Delivery by Doctor</b>	The percentage of live births/deliveries that occur with the assistance of a doctor during the five-year period preceding the survey.
<b>Assisted Delivery by Other Health Professional</b>	The percentage of live births/deliveries that occur with the assistance of a nurse, midwife, village health worker, or any other trained health professional besides a doctor during the five-year period preceding the survey.
<b>Child Survival Indicators</b>	
<b>ARI Treatment - Children Under 5</b>	Percentage of children under five years who were ill with an acute respiratory infection (ARI), which is associated with cough, rapid breathing and a high fever, during the two weeks preceding the survey, and who were treated with specific remedies.
<b>ORT Use Rate (ORS, RHS, or Increased Fluids)</b>	Oral Rehydration Therapy (ORT). Percentage of children under age 5 with diarrhea two weeks prior to survey who received increased fluids, oral rehydration solution (ORS) or recommended home solution (RHS).
<b>Infant Mortality Rate (BUCEN)</b>	The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year.
<b>Infant Mortality Rate (DHS)</b>	The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year (five-year period preceding survey).
<b>Infant Mortality Rate, Females (BUCEN)</b>	The estimated annual number of deaths of female infants under 12 months in a given year per 1,000 live births in that same year.
<b>Infant Mortality Rate, Males (BUCEN)</b>	The estimated annual number of of male infants under 12 months in a given year per 1,000 live births in that same year.
<b>Under-5 Mortality Rate (BUCEN)</b>	Annual number of deaths that occur in children 0-4 years old per 1,000 births.
<b>Under-5 Mortality Rate (DHS)</b>	Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).
<b>Exclusive Breastfeeding (under 4 mos.)</b>	Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.
<b>Exclusive Breastfeeding (under 6 mos.)</b>	Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.
<b>Stunted (height-for-age)</b>	Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.
<b>Underweight (weight-for-age)</b>	Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.
<b>Wasted (weight-for-height)</b>	Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.
<b>DPT3 Vaccination Rate (DHS)</b>	Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.
<b>Measles Vaccination Rate (DHS)</b>	Percentage of living children who have received one dose of MCV (fully immunized) against measles.
<b>Polio Vaccination Rate (DHS)</b>	Percentage of children receiving three doses of polio vaccine.
<b>Tetanus Toxoid Vaccination</b>	Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).

### ***HIV/AIDS Prevention Indicators***

<b>HIV Prevalence, Adults</b>	The estimated number of adults (ages 15-49) living with HIV/AIDS at the end of 1999 divided by the 1999 adult population (ages 15–49).
<b>HIV Prevalence, Females 15-24, High Estimates</b>	HIV prevalence, adolescent females 15-24 for high estimates.
<b>HIV Prevalence, Females 15-24, Low Estimates</b>	HIV prevalence, adolescent females 15-24 for low estimates.
<b>HIV Prevalence, Males 15-24, High Estimates</b>	HIV prevalence, adolescent males 15-24 for high estimates.
<b>HIV Prevalence, Males 15-24, Low Estimates</b>	HIV prevalence, adolescent males 15-24 for low estimates.
<b>Males Reporting Condom Use With Last Non-Regular Partner</b>	Among men who have had sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.
<b>Females Reporting Condom Use With Last Non-Regular Partner</b>	Among women who have had sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.

### ***Infectious Diseases Control Indicators***

<b>TB Incidence Rate</b>	Total number of TB cases (all types) estimated in a given period of time per 100,000 population.
<b>TB Case Detection Rate</b>	TB Case Detection Rate is defined as the percentage of the annual new smear-positive notifications of the estimated annual new smear-positive incidence.
<b>TB (DOTS) Treatment Success Rate</b>	The proportion of smear-positive patients who were cured plus the proportion who completed treatment.
<b>Malaria</b>	Reported number of cases of malaria during the year for which data reported.

# Statistical Sources For Zambia

---

<b>BUCEN - IDB 2000</b>	United States Census Bureau (BUCEN), International Programs Center. International Database. Web site revision dated October 2000.
<b>UNAIDS 2000</b>	Report on the Global HIV/AIDS Epidemic. June 2000. The Joint United Nations Programme on HIV/AIDS, Geneva.
<b>UNESCO 2000</b>	UNESCO Education for All 2000 Assessment. United Nations Educational, Scientific and Cultural Organization, Paris.
<b>WHO/Global Summary 2001</b>	WHO Vaccine-Preventable Diseases Monitoring System. 2001 Global Summary. World Health Organization, Geneva.
<b>WHO/Hill 2001</b>	Kenneth Hill, Carla Abou Zahr & Tessa Wardlaw: Estimates of Maternal Mortality for 1995. Bulletin of the World Health Organization, vol. 79, no. 3, 182-193, 2001.
<b>WHO/RBM 2002</b>	WHO-Roll Back Malaria Web site. A Global Partnership. <a href="http://www.who.int/rbm">http://www.who.int/rbm</a> . Data accessed on web site in March 2002.
<b>World Bank/WDI 2001</b>	World Bank Development Indicators 2001, CD-Rom. Washington, DC: The World Bank, 2001.
<b>Zambia DHS 1992</b>	University of Zambia, Central Statistical Office and Macro International Inc. Zambia Demographic and Health Survey 1992. Columbia, Maryland: Macro International Inc., March 1993.
<b>Zambia DHS 1996</b>	Central Statistics Office [Zambia] and Ministry of Health and Macro International Inc. 1997. "Zambia Demographic and Health Survey, 1996." Calverton, Maryland: Central Statistics Office and Macro International Inc. September 1997.